

Pascoe, King, Macalister, Mitchell. Exhibitors: Crocker, Denham, Simpson, Balls. Hockin Prizeman (for electricity): Browning. Engineering Scholar: Paton.

THE attention of teachers and others engaged in schools is directed to the appeal made by Prof. Karl Pearson in our correspondence columns. Observations of the physical and mental characters of children are required, and measurements of the head, in order to provide material for an investigation of heredity upon which Prof. Pearson is engaged. There should be no difficulty in obtaining the co-operation of masters and mistresses in schools in this work, for the observations and measurements can be made with very little trouble, and they are of as much interest from an educational point of view as they are to biological science.

### SCIENTIFIC SERIAL.

*Bulletin of the American Mathematical Society*, May.—The number opens with four papers read before the Society at the dates annexed: On the geometry of the circle, by Dr. V. Snyder (December 28, 1899); isomorphism between certain systems of single linear groups, by Prof. L. E. Dickson (February 24); the Hessian of the cubic surface ii., by Dr. J. I. Hutchinson (February 24); and note on the group of isomorphisms, by Dr. G. A. Miller (February 24). These papers are short and, in the main, continuations of work previously published by the authors.—Prof. F. S. Woods contributes an interesting sketch of a German translation, by F. Engel, of two articles by Lobachevsky, with the titles "Ueber die Anfangsgründe der Geometrie" and "Neue Anfangsgründe der Geometrie mit einer Vollständigen theorie der Parallellinien." The reviewer's conclusion is that, "while it is remarkable that the solution of a two-thousand-year-old problem should be given almost simultaneously by three men, it should be remembered that these three were not the only mathematicians who had worked upon the problem. More than one had missed the solution by a hair's breadth only; Lobachevsky, Bolyai and Gauss succeeded in finding it."—Other notices are Vogt's "Algebraic solutions of equations," by J. Pierpont; the elements of the differential and integral calculus, based on the work by Nernst and Schönflies (translated by W. A. Young and C. E. Linebarger), by L. E. Dickson; and E. Pascal's "Die Variationsrechnung," by J. K. Whittemore.—University and general mathematical information come into the "Notes" and "New Publications."

### SOCIETIES AND ACADEMIES.

#### LONDON.

**Royal Society**, April 5.—"The Kinetic Theory of Planetary Atmospheres." By Prof. G. H. Bryan, F.R.S.

The application of the kinetic theory to the atmospheres of planets dates from the paper of Waterston, who gave an investigation based on the then only possible assumption of equal velocities for all molecules, an assumption since known as Clausius' law. Of later papers reference is due in especial to Dr. Johnstone Stoney's memoir "Of Atmospheres on Planets and Satellites" (*Trans. R. Dublin Soc.*), in which the test of permanence of a gas in the atmosphere of a planet is made to depend on the ratio of its velocity of mean square to that relative velocity which would enable a suitably projected body to escape from the planet's attraction. If it be admitted, as Dr. Stoney assumes, that helium cannot exist in our atmosphere, it follows that vapour of water cannot exist on Mars.

The author's object has been to investigate the logical conclusions obtained by applying the Boltzmann-Maxwell distribution to the atmospheres of planets. In 1893 calculations were made, having special reference to the absence of atmosphere from the moon, but these took no account of axial rotation. When this cause is taken into account, the distribution of coordinates and relative velocities of the molecules is found to be the same as if the planet were at rest, and "centrifugal force" applied to the system. The surfaces of equal density are of the forms originally investigated by Edward Roche, of Montpellier, and they cease to be closed surfaces when passing to the outside of the point on the equatorial plane where centrifugal force just balances the planet's attraction. Calling the surface through this point the "critical surface," the density of molecular distribution over this surface must be very small to ensure permanence.

NO. 1599, VOL. 62]

The ratio of the density at the planet's surface to the density at the critical surface has been called the "critical density ratio," and the author calculates its logarithm for particular gases at different temperatures on the various planets. The use of this logarithm has the advantage that the calculation can at once be extended to any gas at any temperature.

The high value obtained in the case of helium, considered in reference to the earth, appears to afford abundant proof that if helium existed in our atmosphere it would possess a very high degree of permanence at ordinary temperatures. To test this point further, a calculation is made of the total rate at which molecules would flow across the critical surface, this rate being regarded as a superior limit to the rate at which the planet would lose its atmosphere, since it takes no account of molecules which describe free paths beyond the limit and fall back again. To further exhibit the results in a tangible form, the rate of flow is estimated by the number of years in which the total amount of gas escaping across the critical surface would be equal to the amount of the gas in a layer covering the surface of the planet to the depth of 1 cm. This measure is independent of the actual quantity of the gas under consideration existing in the atmosphere, since, if this quantity be increased, the rate of flow across the critical surface and the amount of gas present in the surface layer 1 cm. thick will be increased in the same proportion.

If a gas of molecular weight 2, such as helium, be supposed to exist in the earth's atmosphere, the loss in question would occupy  $3.5 \times 10^{26}$  years at  $-73^\circ \text{C.}$ ,  $3 \times 10^{19}$  years at  $27^\circ$ ,  $8.4 \times 10^{10}$  years at  $127^\circ \text{C.}$ ,  $6 \times 10^5$  years at  $227^\circ \text{C.}$ , and 222 years at  $327^\circ \text{C.}$

If we have the absolute temperatures, we have the conditions applicable to hydrogen, the losses in question therefore taking place in  $8.4 \times 10^{10}$  years at  $-73^\circ \text{C.}$ ,  $6 \times 10^5$  years at  $-23^\circ \text{C.}$ , and 222 years at  $27^\circ \text{C.}$

For water vapour on Mars, the corresponding results are  $1.2 \times 10^{33}$  years at  $-73^\circ$ ,  $1.9 \times 10^{16}$  years at  $27^\circ$ ,  $2.4 \times 10^9$  years at  $127^\circ$ ,  $4.3 \times 10^5$  years at  $227^\circ$ , and 106 years at  $327^\circ$ .

These figures indicate that helium cannot practically escape from our atmosphere at existing temperatures, nor can vapour of water escape from the atmosphere of Mars. A leakage may, and undoubtedly does, take place, which may appear considerable when estimated by the number of actual molecules escaping, but it is wholly inappreciable relative to the mass of gas left behind.

At a future time it is proposed to examine the corresponding results, based on the hypothesis that the atmosphere of a planet is distributed according to the adiabatic instead of the isothermal law.

"On the Weight of Hydrogen desiccated by Liquid Air." By Lord Rayleigh, F.R.S.

In recent experiments by myself and by others upon the density of hydrogen, the gas has always been dried by means of phosphoric anhydride; and a doubt may remain whether, on one hand, the removal of aqueous vapour is sufficiently complete, and on the other, whether some new impurity may not be introduced. I thought that it would be interesting to weigh hydrogen dried in an entirely different manner, and this I have recently been able to effect with the aid of liquid air, acting as a cooling agent, supplied by the kindness of Prof. Dewar from the Royal Institution. The operations of filling and weighing were carried out in the country as hitherto. I ought, perhaps, to explain that the object was not so much to make a new determination of the highest possible accuracy, as to test whether any serious error could be involved in the use of phosphoric anhydride, such as might explain the departure of the ratio of densities of oxygen and hydrogen from that of 16:1. I may say at once that the result was negative.

Each supply consisted of about six litres of the liquid, contained in two large vacuum-jacketed vessels of Prof. Dewar's design, and it sufficed for two fillings with hydrogen at an interval of two days. The intermediate day was devoted to a weighing of the globe empty. There were four fillings in all, but one proved to be abortive owing to a discrepancy in the weights when the globe was empty, before and after the filling. The gas was exposed to the action of the liquid air during its passage in a slow stream of about half a litre per hour through a tube of thin glass.

I have said that the result was negative. In point of fact the actual weights found were  $\frac{1}{10}$  to  $\frac{2}{10}$  milligrams heavier than in the case of hydrogen dried by phosphoric anhydride. But I